

3-Methylcholanthrene/20-Methylcholanthrene  
(3-MCA)

- 1925      Wieland synthesized 3-MCA from desoxycholic acid. [1]
- 1951      Riegel *et al.* stated "...simultaneous painting of 20-methylcholanthrene and 1,2,5,6-dibenzfluorene delayed skin carcinogenesis in mice beyond the period characteristic of methylcholanthrene alone...." [emphasis added] [2]
- 1961      Stanton and Blackwell induced epidermoid carcinoma in the lungs of rats which received 3-MCA intravenously. [3]
- 1964      Kroller claimed isolation of 3-methylcholanthrene from cigarette smoke; 3-MCA was determined to be carcinogenic to mouse epidermis. [4]
- 1964      Wynder and Hoffmann hypothesized, "since this carcinogenic hydrocarbon has as yet not been found in any other combustion product, it remains a doubtful assumption that it is present in tobacco smoke until analytical criteria such as ultraviolet absorption and fluorescence spectra and further details of isolation method are presented." [5]
- 1976      Dipple, in his article on polynuclear aromatic hydrocarbons concludes, "[E]xperimental studies of the carcinogenic activities of hydrocarbon metabolites indicate that the 3-methylcholanthrene-1-and 2-ones are probably the most active metabolites identified to date." [6]
- 1977      The Royal College of Physicians characterizes polycyclic aromatic hydrocarbons as one of the two (along with N-nitroso compounds) chief initiators of cancer in tobacco smoke.

2501476195

In addition, The Royal College of Physicians reports that "[v]itamin A appears to protect rats against precancerous changes in the living cells and early tumors of the respiratory tract produced by the potent carcinogen 3-methylcholanthrene." [7]

- 1979      The report of the U.S. Surgeon General does not mention 3-methylcholanthrene. [8]
- 1979-      In two separate studies, Rinkus *et al.* (1979) and  
1980      Bartsch *et al.* (1980) found several cigarette smoke constituents to be mutagenic. Among those were 3-MCA. [9]

2501476196

## References

- [1] Wieland, Schlichting, Z. PHYSIOL. CHEM., 150:267 (1925).
  - [2] Riegel *et al.*, CANCER RESEARCH, 11:301 (1951).
  - [3] Stanton, M. F. and Blackwell, R., *Induction of Epidermoid Carcinoma in Lungs of Rats. A "New" Method Based Upon the Deposition of Methylcholanthrene in Areas of Pulmonary Infarction*, J. NAT. CANCER INST., 27:375-407 (1961).
  - [4] Kroller, E., *Comparative Results of Dry Distillation and Smoking of Tobacco*, DEUT. LEBENSM.-RUNDSCHALL, 60:214-215 (1964).
  - [5] Wynder, E. L. and Hoffmann, D., Tobacco and Tobacco Smoke. Studies on Experimental Tobacco Carcinogenesis, 334, Academic Press (1967).
  - [6] Dipple, A. *Polynuclear Aromatic Hydrocarbons*. In: Chemical Carcinogens ACS Monograph 173 Searle, C.E. (Ed.) 245-314 (1976).
  - [7] *Smoking or Health, The Third Report From the Royal College of Physicians of London*, 41 (1977).
  - [8] U.S. Public Health Service, *Smoking and Health: A Report of the Surgeon General* (1979).
  - [9] Rinkus, S.J. and Legator, M.S., *Chemical Characterization of 465 Known or Suspected Carcinogens and Their Correlation with Mutagenic Activity in the Salmonella typhimurium System*, CANCER RES., 39:3289-3318 (1979).
- Bartsch, H., Malaveille, Camus, A.M., Martel-Plauche, G., Brun, G., *et al.*, *Validation and Comparative Studies With S. typhimurium Strains and V79 Chinese Hamster Cells in the Presence of Various Metabolizing Systems*, MUTATION RES., 76:1-50 (1980).

2501476197